



We would like to proudly introduce the finest crosslinked hyaluronic acid dermal filler, Artgaon.

ARTGAON



Company Overview

Artgaon Art for Beauty

Throughout history, science has always made an impact to revolutionize art. We believe that our formula for Artgaon will enable you to create the art you want. Leave the science to us for you to explore the possibilities of beauty imaginable.



Product Lineup

- · Artgaon products contain lidocaine to reduce pain of patients during dermal filler procedures.
- Artgaon Pure products contain no lidocaine for patients who may have allergic reactions to lidocaine.

Joonghun Pharmaceutical



Product Name	Lidocaine	Shelf Life		
Artgaon Shape Plus	0.3%	36 months		
A Concentration	Indicative Target			

20 mg/mL

Needle Size

27 G

Artgaon Shape Plus is indicated to be used for subcutaneous to supraperiosteal injection for cheek augmentation to correct age-related volume defects in mid facial area.



Artgaon Pure Shape	36 months
HA Concentration	Indicative Target
20 mg/mL	Artgaon Pure Shape is indicated to be used for subcutaneous to
Needle Size	supraperiosteal injection for cheek
27 G	augmentation to correct age-related volume defects in mid facial area.

Shelf Life

Product Name

Product Name

Artgaon Pure Deep



Product Name	Lidocaine	Shelf Life	
Artgaon Deep Plus	0.3%	36 months	
HA Concentration	Indicative Target		
20 mg/mL	Artgaon Deep Plus is indicated to be used for mid to deep dermis		
Needle Size	injection for moderate to severe		
27 G	facial wrinkles	and nasolabial folds.	



HA Concentration	Indicative Target
20 mg/mL	Artgaon Pure Deep is indicated to be used for mid to deep dermis
Needle Size	injection for moderate to severe
27 G	facial wrinkles and nasolabial folds.

Shelf Life

36 months



Product Name	Lidocaine	Shelf Life	
Artgaon Fine Plus	0.3%	36 months	
HA Concentration	Indicative Targe	t	
20 mg/mL	Artgaon Fine Plus is indicated to be used for superficial dermis injection		
Needle Size	for fine facial wrinkles and folds or		
30 G	for facial defe	ects.	



Product Name	Shelf Life
Artgaon Pure Fine	36 months
HA Concentration	Indicative Target
20 mg/mL	Artgaon Pure Fine is indicated to be used for superficial dermis injection
Needle Size	for fine facial wrinkles and folds or
30 G	for facial defects.

Artgaon Brochure 04 05

07

Comparison Test Data

Test Categories	Test Specification	Artgon Shape plus	Product of Company X	Product of Company Y
Н	6.0 - 7.5	6.7	N/A	7.0
Osmotic Pressure	200 - 400 mOsm/kg	296 mOsm/kg	304 mOsm/kg	346 mOsm/kg
Injection Force	≤ 40 N	7.2 N (27G)	11.7 N (27G)	14.4 N (27G)
Modulus of Elasticity	N/A	600 Pa	904 Pa	397 Pa
Insoluble Particulate Matter	10 µm : ≤ 6,000 ea/syr 25 µm : ≤ 600 ea/syr	648 ea/syr 30 ea/syr	N/A N/A	8,080 ea/syr 300 ea/syr




Artgaon Brochure 06

Low And Steady Injection Force


Easier for Professionals to Inject and Mold the fillers without damaging the tissue area.

Long Lasting Natural Molding Effect

Injection Force Test




Degradation Test



Artgaon Brochure 08 09

Large & Even Particle Size


Our Particle Size and Shape Controlling Technology provides Long Lasting Natural Look.



Guaranteed Safety for Patients

01. JOONGHUN's technology effectively removes BDDE from the injectable solution.

02. Minimal insoluble particles ensure that no adverse side effects arise.



Reference : 1) Particle(≥10 /m) ≤ 6000ea/syr, Particle(≥25 /m) ≤ 600ea/syr, Korean Pharmacopeia and US Pharmacopeia testing standards.

03. High Purity Indicators.

BDDE Residue ²⁾	LOD
Endotoxin ³⁾	LOD
Biological Safety Test ⁴⁾	Approved in all items


Reference : 2) LOD : 0.1ppm, Joonghun Lab. 3) LOD : 0.005 EU/mL, Joonghun Lab. 4) Korea Testing Institute (KTR)


Artgaon Brochure 10

Clinical Trial⁵⁰ Evaluation Results

The mean values of WSRS and GAIS results for Artgaon showed no significant difference in comparison to the Control Product R.

· Artgaon was proven to be similar to Product R






Joonghun Pharmaceutical

The average value of VAS of Artgaon is significantly low immediately after application of the medical device

· Artgaon was proven to cause less pain than product R immediately after application



Study Design

Single-center, matched-pairs, randomized double blinded pivotal study
Study Period: 24 Weeks · Number of Subjects: 83

Reference Data

9) Clinical study to evaluate the safety and efficacy of Artgaon Shape Plus (vs. Restylane LYFT with Lidocaine) through single-center, prospective, matched-pairs, randomized double blinded pivotal study

Artgaon Brochure 12 13

Contact point

Joonghun Pharmaceutical Co., Ltd. 15 Gukhoe-daero 62-gil, Yeongdeungpo-gu, Seoul, Korea 07236

Tel	82.2.2138.8114
Fax	82.2.6928.9119
E-Mail	artgaon@joonghun.com



Artgaon, the finest for European professionals

Artgaon Brochure 14